

**RESTRICTED USE PESTICIDE**

Due to dermal irritation.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certificate.

**FIFRA Sec. 24(c) Special Local Need  
SLN Label**



**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MINNESOTA**

**SPIN-AID ® HERBICIDE**

**EPA Reg. No. 87865-5  
EPA 24(c) SLN No. MN-230002**

**SPIN-AID will be available for a limited time under a supplemental distribution label with the EPA Reg. No. 264-616-89865. This supplemental distribution label expired on 11/08/2022 and is not legal for use on its own or in conjunction with this 24c label after 05/08/2024.**

**This label for SPIN-AID Herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2027**

**KEEP OUT OF REACH OF CHILDREN**

**BROADCAST APPLICATION POSTEMERGENCE FOR CONTROL OF  
COMMON LAMBSQUARTERS AND KOCHIA IN SUGARBEETS**

**ATTENTION**

- **It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**
- **This labeling must be in the possession of the user at the time of application.**
- **Read the label affixed to the container for SPIN-AID Herbicide before applying.**
- **Carefully follow all precautionary statements and applicable use directions.**
- **Use of SPIN-AID Herbicide according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for SPIN-AID Herbicide.**

## SPECIFIC USE DIRECTIONS

Refer to product label for Mixing, Handling and Application Instructions.

### RATES OF APPLICATION – SUGARBEET

When applying the following recommendations for SPIN-AID® Herbicide to sugarbeets, add HSMOC at 1.0 to 1.5pt/A.

**By Ground:** Multiple applications of SPIN-AID® Herbicide may be applied by ground to sugarbeets to control common lambsquarters and kochia. Broadcast Application rates are provided in appropriate charts in this section. The first application MUST NOT be applied before the sugarbeets have reached the 2-leaf stage. The Spin-Aid rate is adjusted for sugarbeet crop stage to ensure crop safety. For broadcast applications with ground equipment, apply in 20 to 30 gallons of water per acre. To completely control the targeted weeds, a second application timed to 5 to 7 days after the first application is REQUIRED. A maximum of three (3) applications are allowed and not to exceed a total of 3 qts/A.

For further information, contact Belchim Crop Protection USA: [info.usa@belchimusa.com](mailto:info.usa@belchimusa.com)

### USE PRECAUTIONS

DO NOT APPLY SPIN-AID® HERBICIDE TO SUGARBEETS LATER THAN 75 DAYS PRIOR TO HARVEST.

For sugarbeets, leave a 16 foot buffer from the application area when the wind direction is toward sensitive dicot plants.

### SPIN-AID® HERBICIDE DOSAGE CHART FOR BROADCAST APPLICATION - SUGARBEETS

**Common Lambsquarters Control Programs: Rates to be applied based upon crop stage when weeds emerge**

Two Application Program	Application Number	Sugarbeet Stage	Spin-Aid + Glyphosate (Fl oz/A + lb ai/A)	OR	Spin-Aid + ethofumesate + Glyphosate (Fl oz/A + lb ai/A)
	1	2-leaf	16oz + 1.15lb		12oz + 4oz + 1.15lb
Again 5-7 days later	2	4-leaf	32oz + 1.15lb		24oz + 4oz + 1.15lb
OR					
	1	4-leaf	32oz + 1.15lb		24oz + 4oz + 1.15lb
Again 5-7 days later	2	6-leaf	32oz + 1.15lb		24oz + 4oz + 1.15lb
Add HSMOC at 1.0 to 1.5 pt/A in 20-30 GPA water to all tank mixtures					

**Kochia Control Programs: DIME SIZE – 1 inch**

Two Application Program	Application Number	Sugarbeet Stage	Spin-Aid + Glyphosate (Fl oz/A + lb ai/A)	OR	Spin-Aid + ethofumesate + Glyphosate (Fl oz/A + lb ai/A)
	1	2-leaf	24oz + 1.15lb		16oz + 4oz + 1.15lb
Again 5-7 days later	2	4-leaf	32oz + 1.15lb		24oz + 4oz + 1.15lb
OR					
	1	4-leaf	32oz + 1.15lb		24oz + 4oz + 1.15lb
Again 5-7 days later	2	6-leaf	40oz + 1.15lb		32oz + 4oz + 1.15lb
Add HSMOC at 1.0 to 1.5 pt/A in 20-30 GPA water to all tank mixtures					

**Kochia Control Programs: QUARTER SIZE – 2 inch**

Two Application Program	Application Number	Sugarbeet Stage	Spin-Aid + Glyphosate (Fl oz/A + lb ai/A)	OR	Spin-Aid + ethofumesate + Glyphosate (Fl oz/A + lb ai/A)
	1	2-leaf	24oz + 1.15lb		16oz + 4oz + 1.15lb
Again 5-7 days later	2	4-leaf	36oz + 1.15lb		32oz + 4oz + 1.15lb
OR					
	1	4-leaf	36oz + 1.15lb		32oz + 4oz + 1.15lb
Again 5-7 days later	2	6-leaf	48oz + 1.15lb		48oz + 4oz + 1.15lb
Add HSMOC at 1.0 to 1.5 pt/A in 20-30 GPA water to all tank mixtures					