

Herbicide

Spin-Aid®

**Post Emergent Herbicide for
Control of Weeds in Spinach
(Processing & Seed Only)
and Red (Garden) Beets**

Active Ingredient:

Phenmedipham* 15.9%

Spin-Aid® is an extremely effective post emergent herbicide for control of broad leaf weeds in spinach (grown for processing and seeding) as well as red garden beets. Belchim Crop Protection is pleased to be the exclusive marketer of this very effective post emergent herbicide, keeping the product available to the industry.

Spin-Aid can be applied to young processing spinach and red beets for the control of several key broad leaf weeds.

Wild mustard ... *Brassica kaber*
Coast fiddleneck ... *Amsinckia intermedia*
Common lambsquarters ... *Chenopodium album*
Common chickweed ... *Stellaria media*
Shepherdspurse *Capsella bursa-pastoris*
Purslane ... *Portulaca oleracea*
London rocket ... *Sisymbrium irio*
Common ragweed *Ambrosia artemisiifolia*
Nettleleaf goosefoot ... *Chenopodium murale*
Annual sowthistle ... *Sonchus oleraceus*
Groundcherry ... *Physalis lanceifolia*

* 3-methoxycarbonylaminophenyl-3- methylcarbanilate
Contains 1.3 lbs. active ingredient per gallon.

Note: Spin-Aid is not registered in all states.

Contact your Belchim Crop Protection representative for the most up to date information.

Read entire product label prior to use for complete use instructions.

Application Methods:

Spinach

Ground: Apply at the rate of 3 to 6 pints per acre to spinach at the 4-true leaf stage or larger in 10 to 20 gallons of water on a broadcast basis.

Band application: See label for dosage chart.

Air: Apply at the rate of 3 to 6 pints per acre to spinach at the 4-true leaf stage or larger using 5 to 20 gallons of spray per acre.

Reduced Rates: Can be applied as a split application for weed control. The first application of only 2 1/2 to 3 pints per acre can be applied at the 2 to 4-leaf stage of spinach (instead of the 4 to 6 true-leaf stage) followed by a sequential application of only 3 pints, 4 to 6 days later. Use maximum of 6 pints per acre per year total with this method.

Red (Garden) Beets

Ground: Multiple applications may be applied by ground to red (garden) beets to control early germinating weeds. See Label for Broadcast and Band Application rates. The first application must be applied when the red (garden) beets have reached the 2-leaf stage. Any weeds which are not completely controlled by the first treatment will usually be checked or controlled by repeat applications.

Contact your BELCHIM CROP PROTECTION representative or a local extension agent for additional information and application recommendations.

Packaging:

2 x 2.5 gallon jug
36 cases per pallet

Marketed by:



BELCHIM
CROP PROTECTION

2751 Centerville Road | Suite 100
Wilmington, DE 19808 | Phone: 855-445-7990
www.belchimusa.com | info.usa@belchim.com