

Belchim Crop Protection US Corporation 2751 Centerville Road, Suite 100 Wilmington, DE 19808 855-445-7990 www.belchimusa.com

FIFRA 2(ee) Recommendation FOR THE USE OF SPIN-AID HERBICIDE FOR POSTEMERGENCE CONTROL OF WEEDS IN SPINACH (GROWN FOR PROCESSING AND SEED)

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin must be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARIZONA AND CALIFORNIA.

Spin-Aid Herbicide EPA Reg. No. 264-616-87865

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

CROP	WEEDS	RATE/ACR E	USE RECOMMENDATIONS
SPINACH	Wild mustard Brassica kaber Common lambsquarters Chenopodium album Shepherdspurse Capsella bursa-pastoris London rocket Sisymbrium irio Nettleleaf goosefoot Chenopodium murale Groundcherry Physalis lanceifolia Coast fiddleneck Amsinckia intermedia Common chickweed Stellaria media Purslane Portulaca oleracea Common ragweed Ambrosia artemisiifolia Annual sowthistle Sonchus oleraceus	Up To 1 Pint/Acre	 Apply 1 pint of Spin-Aid 7 -10 days after seeding when weeds are at two- leaf stage. Note: This application should follow a pre emergent application of Cycloate Herbicide Please refer to USE PRECAUTIONS on Label for suggested application climatic conditions

For Chemical Emergency, Spill, Leak, Fire, Exposure or Accident, Call CHEMTREC Day or Night Within USA and Canada 1-800-424-9300

Spin-Aid[®] is manufactured by and is a registered trademark of:

Bayer CropScience